

A New Numerical Procedure to Compute the Residues of a Complex Functions

Abbas Y. Al-Bayati

profabbasalbayati@yahoo.com

College of Computer Sciences
and Mathematics

University of Mosul/Iraq

Sasan A. Al-Shwani

College of Science

University of Kirkuk

Received on: 27/08/2006

Accepted on: 16/11/2006

ABSTRACT

In this paper, we are going to deal with computations of **Residues and Poles** for the complex functions . We are also going to investigate a **new** numerical procedure theoretically and its implementation numerically to compute the **residue** of complex analytic functions with high order poles. The paper needs the knowledge of computing the complex improper integrations.

Keywords: complex analytic functions, Residues and Poles, complex improper integrations.

طريقة عددية لحساب الرواسب والأقطاب في الدوال العقدية

عباس يونس البياتي

كلية علوم الحاسوب والرياضيات/جامعة الموصل

تاريخ قبول البحث: 2006/11/16

ساسان عادل الشواني

جامعة كركوك/كلية العلوم

تاريخ استلام البحث: 2006/8/27

المخلص

في هذا البحث سوف نتطرق إلى حساب الرواسب والأقطاب في الدوال العقدية . كذلك سوف نحاول التقصي في استحداث وسيلة عددية جديدة نظريا وعمليا لحساب الرواسب للدوال العقدية القابلة للتحليل للأقطاب ذات الرتب العالية. ويحتاج هذا البحث إلى معرفة حسابات التكاملات العقدية المعتلة.

الكلمات المفتاحية: دوال عقدية قابلة للتحليل، الرواسب والاقطاب، التكاملات العقدية المعتلة.

1-Introduction :

A final bit of theory in this paper is devoted to Cauchy's residue theorem and its applications . Recall that Cauchy's theorem stated that ($\oint f(z)dz = 0$) so long as $f(z)$ was analytic every where inside C . What if $f(z)$ is not analytic within C ?

The answer is provided by Cauchy's integral formula . Here we have proposed a **new** procedure to compute the complex improper integrations by calculating the **residues** of the function .

To compute the residues of a complex function of the form $f(z) = \frac{p(z)}{q(z)}$ such that the function $f(z)$ has a pole at $(z = z_0)$, we are going to use the (short – cut - method) to estimate the residues[9].

If R is the real field and C is the complex field then consider the following definitions and theorems:

1-1 poles :[3]

If we find a positive integer (m) such that $[\lim_{z \rightarrow z_0} (z - z_0)^m f(z) \neq 0]$ Then $(z = z_0)$ is called a pole of order (m) . If $(m=1)$ (z_0) is called a simple pole.

1-2 Analytic functions :[9]

If the derivative $f'(z)$ exists at all points (z) of a region R ,then $f(z)$ is said to be analytic in R .

1-3 Taylor's theorem :[3]

Let $f(z)$ be analytic inside and on a simple closed curve C . Let (a) and $(a+h)$ be two points inside C , then

$$f(a+h) = f(a) + h f'(a) + \frac{h^2}{2!} f''(a) + \dots \quad \text{or writing as}$$

$$f(z) = f(a) + (z-a) f'(a) + \frac{(z-a)^2}{2!} f''(a) + \dots$$

Where $(z=a+h)$

The above series is called Taylor expansion for $f(z)$.

1-4 Cauchy integral formula :[2]

If $f(z)$ is analytic inside and on a simple closed curve C and (a) is any point inside C then

$$f(a) = \frac{1}{2\pi i} \int_c \frac{f(z)}{(z-a)} dz$$

Where C is traversed in the positive sense also the n th derivative of $(z=a)$ is given by

$$f^{(n)}(a) = \frac{1}{2\pi i} \int_c \frac{f(z)}{(z-a)^{n+1}} dz \quad ; \quad n=1,2,3,\dots$$

1-5 Lemma (1) :[9]

Let (Z_o) be a pole of order (m) of a function $f(z)$ then the residue of the function at a pole given by the form :

$$\text{Res}(f, Z_o) = \frac{1}{(m-1)!} \lim_{z \rightarrow z_o} \frac{d^{m-1}}{dz^{m-1}} [(z - z_o)^m f(z)].$$

The formula namely (short – cut – method) .

2- A New method to compute the residues of a complex function:

If $f(z)$ has a pole $(z = z_o)$ of order (m) then

$$\frac{1}{2\pi i} \oint_c f(z) dz = \frac{1}{(m-1)!} \lim_{z \rightarrow z_o} \frac{d^{m-1}}{dz^{m-1}} [(z - z_o)^m f(z)] \quad \text{where} \quad [f(z) = \frac{p(z)}{q(z)}]$$

We know that :

$$\text{Res}(f, z_o) = \frac{1}{(m-1)!} \lim_{z \rightarrow z_o} \frac{d^{m-1}}{dz^{m-1}} [(z - z_o)^m f(z)]$$

We find the value of the above integration by calculating the residues by this procedure and it can be found in [5].

2.1) If $(z = z_o)$ is a simple pole i.e $[q(z) = 0, q'(z_o) \neq 0]$ then the residue is given by the following form :

$$\text{Res}(f, z_o) = \frac{p(z_o)}{q'(z_o)} \quad \dots (1)$$

2.2) If $(z = z_o)$ is a pole of order $(m=2)$ i.e $[q(z_o) = 0, q'(z_o) = 0, q''(z_o) \neq 0]$ then the residue is given by the following form :

$$\text{Res}(f, z_o) = 2 \times \frac{p'(z_o)}{q^{(2)}(z_o)} - \frac{2}{3} \times \frac{p(z_o) \cdot q^{(3)}(z_o)}{(q^{(2)}(z_o))^2} \quad \dots (2)$$

See [5] for the details of these procedures.

3- In this paper we are going to follow a procedure for a general function of a pole of order (m):

3.1) Let us start with order (m=3):

If $(z = z_o)$ is a pole of order $(m=3)$ i.e

$[q(z_o) = 0, q'(z_o) = 0, q''(z_o) = 0, q'''(z_o) \neq 0]$ then the residue is given by the following form :

$$\begin{aligned} \operatorname{Res}(f, z_0) = & 3 \times \left[\frac{p^{(2)}(z_0)}{q^{(3)}(z_0)} - \frac{1}{10} \times \frac{p(z_0) \cdot q^{(5)}(z_0)}{(q^{(3)}(z_0))^2} \right. \\ & \left. - \frac{1}{2} \times \frac{q^{(4)}(z_0) p^{(1)}(z_0)}{(q^{(3)}(z_0))^2} + \frac{1}{8} \times \frac{(q^{(4)}(z_0))^2 p(z_0)}{(q^{(3)}(z_0))^3} \right] \end{aligned} \quad \dots(3)$$

Proof:

If $(z = z_0)$ is a pole of order $(m=3)$ then by (short-cut-method) we get :

$$\operatorname{Res}(f, z_0) = \frac{1}{2!} \lim_{z \rightarrow z_0} \frac{d^2}{dz^2} \left[(z - z_0)^3 \frac{p(z)}{q(z)} \right] \quad \dots\dots\dots(4)$$

We expand the analytic function $q(z)$ into Taylor series valid in disk $|z - z_0| < r$

$$= \frac{1}{2} \lim_{z \rightarrow z_0} \frac{d^2}{dz^2} \left[\frac{p(z)(z - z_0)^3}{q(z_0) + (z - z_0)q^{(1)}(z_0) + \frac{(z - z_0)^2}{2!} q^{(2)}(z_0) + \dots\dots\dots} \right] \quad \dots\dots\dots(5)$$

$$= 3 \lim_{z \rightarrow z_0} \frac{d^2}{dz^2} \left[\frac{p(z)}{U(z)} \right] \quad \dots\dots\dots(6)$$

Where

$$U(z) = q^{(3)}(z_0) + \frac{(z - z_0)}{4} q^{(4)}(z_0) + \frac{(z - z_0)^2}{5 \times 4} q^{(5)}(z_0) + \frac{(z - z_0)^3}{6 \times 5 \times 4} q^{(6)}(z_0) + \dots \quad \dots\dots(7)$$

Then from (6) we get :

$$= 3 \lim_{z \rightarrow z_0} \frac{d}{dz} \left[\frac{U(z) p^{(1)}(z) - p(z) U^{(1)}(z)}{(U(z))^2} \right] \quad \dots\dots\dots(8)$$

$$\begin{aligned} = & 3 \lim_{z \rightarrow z_0} \left[\frac{U(z) p^{(2)}(z) - p^{(1)}(z) U^{(1)}(z)}{(U(z))^2} \right. \\ & \left. - \frac{p(z) U^{(2)}(z) + U^{(1)}(z) p^{(1)}(z)}{(U(z))^2} - 2 \times \frac{(U^{(1)}(z))^2 p(z)}{(U(z))^3} \right] \end{aligned} \quad \dots\dots\dots(9)$$

$$\left. \begin{aligned}
 \lim_{z \rightarrow z_0} p(z) &= p(z_0) \\
 \lim_{z \rightarrow z_0} p^{(1)}(z) &= p^{(1)}(z_0) \\
 U(z) &= q^{(3)}(z_0) + \frac{(z-z_0)}{4} q^{(4)}(z_0) + \frac{(z-z_0)^2}{5 \times 4} q^{(5)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U(z) &= q^{(3)}(z_0) \\
 U^{(1)}(z) &= \frac{1}{4} q^{(4)}(z_0) + \frac{2}{5 \times 4} (z-z_0) q^{(5)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U^{(1)}(z) &= \frac{1}{4} q^{(4)}(z_0) \\
 U^{(2)}(z) &= \frac{2}{5 \times 4} q^{(5)}(z_0) + \frac{3 \times 2}{6 \times 5 \times 4} (z-z_0) q^{(6)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U^{(2)}(z) &= \frac{1}{10} q^{(5)}(z_0)
 \end{aligned} \right\} \dots\dots(10)$$

Then Substituting the above values in (10) in equation (9) we get the desired proof .

3.2) If $(z = z_0)$ is a pole of order **(m=4)** i.e

$[q(z_0) = 0, q'(z_0) = 0, q''(z_0) = 0, q'''(z_0) = 0, q^{(4)}(z_0) \neq 0]$ then the residue is given by the following form :

$$\begin{aligned}
 \text{Res}(f, z_0) &= 4 \times \left[\frac{p^{(3)}(z_0)}{q^{(4)}(z_0)} - \frac{3}{5} \times \frac{p^{(2)}(z_0)q^{(5)}(z_0)}{(q^{(4)}(z_0))^2} \right. \\
 &\quad - \frac{1}{35} \times \frac{p(z_0)q^{(7)}(z_0)}{(q^{(4)}(z_0))^2} - \frac{1}{5} \times \frac{p^{(1)}(z_0)q^{(6)}(z_0)}{(q^{(4)}(z_0))^2} \\
 &\quad + \frac{2}{25} \times \frac{p(z_0)q^{(5)}(z_0)q^{(6)}(z_0)}{(q^{(4)}(z_0))^3} + \frac{6}{25} \times \frac{(q^{(5)}(z_0))^2 p^{(1)}(z_0)}{(q^{(4)}(z_0))^3} \\
 &\quad \left. - \frac{6}{125} \times \frac{(q^{(5)}(z_0))^3 p(z_0)}{(q^{(4)}(z_0))^4} \right] \dots\dots(11)
 \end{aligned}$$

Proof:

If $(z = z_0)$ is a pole of order **(m=4)** then by (short – cut - method) we get :

$$\operatorname{Res}(f, z_0) = \frac{1}{3!} \lim_{z \rightarrow z_0} \frac{d^3}{dz^3} \left[(z - z_0)^4 \frac{p(z)}{q(z)} \right] \quad \dots\dots(12)$$

We expand the analytic function $q(z)$ into Taylor series valid in disk $|z - z_0| < r$

$$= \frac{1}{3!} \lim_{z \rightarrow z_0} \frac{d^3}{dz^3} \left[\frac{(z - z_0)^4 p(z)}{q(z_0) + (z - z_0)q^{(1)}(z_0) + \frac{(z - z_0)^2}{2!} q^{(2)}(z_0) + \dots} \right] \quad \dots\dots(13)$$

$$= 4 \lim_{z \rightarrow z_0} \frac{d^3}{dz^3} \left[\frac{p(z)}{q(z)} \right] \quad \dots\dots(14)$$

Where :

$$U(z) = q^{(4)}(z_0) + \frac{(z - z_0)}{5} q^{(5)}(z_0) + \frac{(z - z_0)^2}{6 \times 5} q^{(6)}(z_0) + \dots \quad \dots\dots(15)$$

Then from (14) we get :

$$= 4 \lim_{z \rightarrow z_0} \frac{d^2}{dz^2} \left[\frac{U(z) p^{(1)}(z) - p(z) U^{(1)}(z)}{(U(z))^2} \right] \quad \dots\dots\dots(16)$$

$$= 4 \lim_{z \rightarrow z_0} \frac{d}{dz} \left[\frac{U(z) p^{(2)}(z) - p^{(1)}(z) U^{(1)}(z)}{(U(z))^2} - \frac{p(z) U^{(2)}(z) + U^{(1)}(z) p^{(1)}(z)}{(U(z))^2} - 2 \times \frac{(U^{(1)}(z))^2 p(z)}{(U(z))^3} \right] \quad \dots\dots\dots(17)$$

$$= 4 \lim_{z \rightarrow z_0} \left[\frac{p^{(3)}(z)}{U(z)} - 3 \times \frac{p^{(2)}(z) U^{(1)}(z)}{(U(z))^2} - \frac{p(z) U^{(3)}(z)}{(U(z))^2} - 3 \times \frac{p^{(1)}(z) U^{(2)}(z)}{(U(z))^2} + 6 \times \frac{p(z) U^{(1)}(z) U^{(2)}(z)}{(U(z))^3} + 6 \times \frac{(U^{(1)}(z))^2 p^{(1)}(z)}{(U(z))^3} - 6 \times \frac{(U^{(1)}(z))^3 p(z)}{(U(z))^4} \right] \quad \dots\dots(18)$$

Where

$$\left. \begin{aligned} U(z) &= q^{(4)}(z_0) + \frac{1}{5}(z - z_0)q^{(5)}(z_0) + \frac{1}{6 \times 5}(z - z_0)^2 q^{(6)}(z_0) + \dots ; \lim_{z \rightarrow z_0} U(z) = q^{(4)}(z_0) \\ U^{(1)}(z) &= \frac{1}{5}q^{(5)}(z_0) + \frac{2}{6 \times 5}(z - z_0)q^{(6)}(z_0) + \dots ; \lim_{z \rightarrow z_0} U^{(1)}(z) = \frac{1}{5}q^{(5)}(z_0) \\ U^{(2)}(z) &= \frac{2}{6 \times 5}q^{(6)}(z_0) + \frac{6}{7 \times 6 \times 5}(z - z_0)q^{(7)}(z_0) + \dots ; \lim_{z \rightarrow z_0} U^{(2)}(z) = \frac{1}{15}q^{(6)}(z_0) \\ U^{(3)}(z) &= \frac{6}{7 \times 6 \times 5}q^{(7)}(z_0) + \frac{24}{8 \times 7 \times 6 \times 5}(z - z_0)q^{(8)}(z_0) + \dots ; \lim_{z \rightarrow z_0} U^{(3)}(z) = \frac{1}{35}q^{(7)}(z_0) \end{aligned} \right\} \quad \dots\dots(19)$$

Then substituting the above values in (19) in equation (18) we get the desired proof.

3.3) In this approach we can find the residue for (m=5):

If $(z = z_o)$ is a pole of order (m=5) i.e

$[q(z_o) = 0, q'(z_o) = 0, q''(z_o) = 0, q'''(z_o) = 0, q^{(4)}(z_o) = 0, q^{(5)}(z_o) \neq 0]$ then the residue is given by the following form :

$$\begin{aligned}
 \text{Res}(f, z_o) = & 5 \times \left[\frac{p^{(4)}(z_o)}{q^{(5)}(z_o)} - 4 \times \frac{\frac{1}{6} q^{(6)}(z_o) p^{(3)}(z_o)}{(q^{(5)}(z_o))^2} \right. \\
 & - 6 \times \frac{\frac{1}{21} q^{(7)}(z_o) p^{(2)}(z_o)}{(q^{(5)}(z_o))^2} + 12 \times \frac{(\frac{1}{6} q^{(6)}(z_o))^2 p^2(z_o)}{(q^{(5)}(z_o))^3} \\
 & - \frac{\frac{1}{126} q^{(9)}(z_o) p(z_o)}{(q^{(5)}(z_o))^2} - 4 \times \frac{\frac{1}{56} q^{(8)}(z_o) p^{(1)}(z_o)}{(q^{(5)}(z_o))^2} \\
 & + \frac{\frac{1}{42} q^{(6)}(z_o) q^{(8)}(z_o) p(z_o)}{(q^{(5)}(z_o))^3} \\
 & + 4 \times \frac{\frac{1}{21} q^{(6)}(z_o) q^{(7)}(z_o) p^{(1)}(z_o)}{(q^{(5)}(z_o))^3} + 6 \times \frac{(\frac{1}{21} q^{(7)}(z_o))^2 p(z_o)}{(q^{(5)}(z_o))^3} \\
 & - 36 \times \frac{(\frac{1}{6} q^{(6)}(z_o))^2 \frac{1}{21} q^{(7)}(z_o) p(z_o)}{(q^{(5)}(z_o))^4} \\
 & \left. - 24 \times \frac{(\frac{1}{6} q^{(6)}(z_o))^3 p^{(1)}(z_o)}{(q^{(5)}(z_o))^4} + 24 \times \frac{(\frac{1}{6} q^{(6)}(z_o))^4 p(z_o)}{(q^{(5)}(z_o))^5} \right] \dots(20)
 \end{aligned}$$

Proof : (By the same way in (m=3)&(m=4)) .

3.4) In this approach we can find the residue for (m=6):

If $(z = z_o)$ is a pole of order (m=6) i.e

$[q(z_o) = q^{(1)}(z_o) = q^{(2)}(z_o) = q^{(3)}(z_o) = q^{(4)}(z_o) = q^{(5)}(z_o) = 0, q^{(6)}(z_o) \neq 0]$ then the residue is given by the following form :

$$\begin{aligned}
 \text{Res}(f, z_0) = & 6 \times \left[\frac{p^{(5)}(z_0)}{q^{(6)}(z_0)} - 5 \times \frac{\frac{1}{7} p^{(4)}(z_0) q^{(7)}(z_0)}{(q^{(6)}(z_0))^2} \right. \\
 & - 4 \times \frac{\frac{1}{28} p^{(3)}(z_0) q^{(8)}(z_0)}{(q^{(6)}(z_0))^2} - 10 \times \frac{\frac{1}{84} p^{(2)}(z_0) q^{(9)}(z_0)}{(q^{(6)}(z_0))^2} \\
 & - 6 \times \frac{\frac{1}{28} p^{(2)}(z_0) q^{(8)}(z_0)}{(q^{(6)}(z_0))^2} - 5 \times \frac{\frac{1}{210} p^{(1)}(z_0) q^{(10)}(z_0)}{(q^{(6)}(z_0))^2} \\
 & - \frac{\frac{1}{462} p(z_0) q^{(11)}(z_0)}{(q^{(6)}(z_0))^2} + 20 \times \frac{(\frac{1}{7} q^{(7)}(z_0))^2 p^{(3)}(z_0)}{(q^{(6)}(z_0))^3} \\
 & + 60 \times \frac{\frac{1}{196} q^{(7)}(z_0) q^{(8)}(z_0) p^{(2)}(z_0)}{(q^{(6)}(z_0))^3} + 10 \times \frac{(\frac{1}{7} \times \frac{1}{210}) p(z_0) q^{(7)}(z_0) q^{(10)}(z_0)}{(q^{(6)}(z_0))^3} \\
 & + 8 \times \frac{\frac{1}{196} q^{(7)}(z_0) q^{(8)}(z_0) p^{(1)}(z_0)}{(q^{(6)}(z_0))^3} + 20 \times \frac{(\frac{1}{84} \times \frac{1}{28}) q^{(9)}(z_0) q^{(8)}(z_0) p(z_0)}{(q^{(6)}(z_0))^3} \\
 & + 32 \times \frac{(\frac{1}{196}) q^{(7)}(z_0) q^{(9)}(z_0) p^{(1)}(z_0)}{(q^{(6)}(z_0))^3} + 24 \times \frac{\frac{1}{196} p^{(1)}(z_0) q^{(7)}(z_0) q^{(9)}(z_0)}{(q^{(6)}(z_0))^3} \\
 & + 30 \times \frac{(\frac{1}{28} q^{(8)}(z_0))^2 p^{(1)}(z_0)}{(q^{(6)}(z_0))^3} - 60 \times \frac{(\frac{1}{7} q^{(7)}(z_0))^3 p^{(2)}(z_0)}{(q^{(6)}(z_0))^4} \\
 & - 60 \times \frac{(\frac{1}{7} q^{(7)}(z_0))^2 \frac{1}{84} q^{(9)}(z_0) p(z_0)}{(q^{(6)}(z_0))^4} - 180 \times \frac{(\frac{1}{7} q^{(7)}(z_0))^2 \frac{1}{28} q^{(8)}(z_0) p^{(1)}(z_0)}{(q^{(6)}(z_0))^4} \\
 & - 90 \times \frac{\frac{1}{7} q^{(7)}(z_0) (\frac{1}{28} q^{(8)}(z_0))^2 p(z_0)}{(q^{(6)}(z_0))^4} + 240 \times \frac{(\frac{1}{7} q^{(7)}(z_0))^3 \frac{1}{28} q^{(8)}(z_0) p(z_0)}{(q^{(6)}(z_0))^5} \\
 & \left. + 120 \times \frac{(\frac{1}{7} q^{(7)}(z_0))^4 p^{(1)}(z_0)}{(q^{(6)}(z_0))^5} - 120 \times \frac{(\frac{1}{7} q^{(7)}(z_0))^5 p(z_0)}{(q^{(6)}(z_0))^6} \right] \dots(21)
 \end{aligned}$$

Proof: (By the same way in (m=3)&(m=4)) .

3.5 If $(z = z_0)$ is a pole of order (m=7) i.e

$$\left[q(z_0) = q^{(1)}(z_0) = q^{(2)}(z_0) = q^{(3)}(z_0) = q^{(4)}(z_0) = q^{(5)}(z_0) = q^{(6)}(z_0) = 0, q^{(7)}(z_0) \neq 0 \right]$$

then the residue is given by the following form:

$$\begin{aligned}
 \text{Res}(f, z) = & 7 \times \left[\frac{p^{(6)}(z_o)}{q^{(7)}(z_o)} - 6 \times \frac{p^{(5)}(z_o) \frac{1}{8} q^{(8)}(z_o)}{(q^{(7)}(z_o))^2} \right. \\
 & - 9 \times \frac{p^{(4)}(z_o) \frac{1}{36} q^{(9)}(z_o)}{(q^{(7)}(z_o))^2} - 14 \times \frac{p^{(3)}(z_o) \frac{1}{120} q^{(10)}(z_o)}{(q^{(7)}(z_o))^2} - \\
 & - 15 \times \frac{\frac{1}{330} q^{(11)}(z_o) p^{(2)}(z_o)}{(q^{(7)}(z_o))^2} - 6 \times \frac{p^{(2)}(z_o) \frac{1}{120} q^{(10)}(z_o)}{(q^{(7)}(z_o))^2} \\
 & - 6 \times \frac{p^{(3)}(z_o) \frac{1}{36} q^{(9)}(z_o)}{(q^{(7)}(z_o))^2} - 6 \times \frac{p^{(1)}(z_o) \frac{1}{792} q^{(12)}(z_o)}{(q^{(7)}(z_o))^2} - \\
 & \frac{p(z_o) \frac{1}{1716} q^{(13)}(z_o)}{(q^{(7)}(z_o))^2} + 20 \times \frac{(\frac{1}{8} \times \frac{1}{36}) q^{(8)}(z_o) q^{(9)}(z_o) p^{(2)}(z_o)}{(q^{(7)}(z_o))^3} \\
 & + 30 \times \frac{(\frac{1}{8} q^{(8)}(z_o))^2 p^{(4)}(z_o)}{(q^{(7)}(z_o))^3} + 108 \times \frac{(\frac{1}{8} \times \frac{1}{36}) q^{(8)}(z_o) q^{(9)}(z_o) p^{(3)}(z_o)}{(q^{(7)}(z_o))^3} \\
 & + 80 \times \frac{(\frac{1}{8} \times \frac{1}{120}) q^{(8)}(z_o) q^{(10)}(z_o) p^{(2)}(z_o)}{(q^{(7)}(z_o))^3} + 76 \times \frac{(\frac{1}{8} \times \frac{1}{330}) q^{(8)}(z_o) q^{(11)}(z_o) p^{(1)}(z_o)}{(q^{(7)}(z_o))^3} \\
 & + 12 \times \frac{(\frac{1}{8} \times \frac{1}{792}) q^{(8)}(z_o) q^{(12)}(z_o) p(z_o)}{(q^{(7)}(z_o))^3} + 90 \times \frac{(\frac{1}{36} q^{(9)}(z_o))^2 p^{(2)}(z_o)}{(q^{(7)}(z_o))^3} + \\
 & + 30 \times \frac{(\frac{1}{36} \times \frac{1}{330}) p(z_o) q^{(11)}(z_o) q^{(9)}(z_o)}{(q^{(7)}(z_o))^3} + 8 \times \frac{(\frac{1}{8} \times \frac{1}{120}) q^{(8)}(z_o) q^{(10)}(z_o) p^{(1)}(z_o)}{(q^{(7)}(z_o))^3} \\
 & + 8 \times \frac{(\frac{1}{36} q^{(9)}(z_o))^2 p^{(1)}(z_o)}{(q^{(7)}(z_o))^3} + 136 \times \frac{(\frac{1}{36} \times \frac{1}{120}) q^{(9)}(z_o) q^{(10)}(z_o) p^{(1)}(z_o)}{(q^{(7)}(z_o))^3} \\
 & + 20 \times \frac{(\frac{1}{120} q^{(10)}(z_o))^2 p(z_o)}{(q^{(7)}(z_o))^3} + 56 \times \frac{(\frac{1}{8} \times \frac{1}{120}) q^{(8)}(z_o) q^{(10)}(z_o) p^{(2)}(z_o)}{(q^{(7)}(z_o))^3}
 \end{aligned}$$

$$\begin{aligned}
 & -120 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^3 p^{(3)}(z_o)}{(q^{(7)}(z_o))^4} - 540 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^2 \frac{1}{36} q^{(9)}(z_o) p^{(2)}(z_o)}{(q^{(7)}(z_o))^4} \\
 & -90 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^2 \frac{1}{330} q^{(11)}(z_o) p(z_o)}{(q^{(7)}(z_o))^4} - 24 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^2 \frac{1}{36} q^{(9)}(z_o) p^{(1)}(z_o)}{(q^{(7)}(z_o))^4} \\
 & -360 \times \frac{\left(\frac{1}{8} \times \frac{1}{36} \times \frac{1}{120}\right) q^{(8)}(z_o) q^{(9)}(z_o) q^{(10)}(z_o) p(z_o)}{(q^{(7)}(z_o))^4} - \\
 & -540 \times \frac{\left(\frac{1}{36} q^{(9)}(z_o)\right)^2 \frac{1}{8} q^{(8)}(z_o) p^{(1)}(z_o)}{(q^{(7)}(z_o))^4} - 90 \times \frac{\left(\frac{1}{36} q^{(9)}(z_o)\right)^2 p(z_o)}{(q^{(7)}(z_o))^4} \\
 & + 360 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^4 p^{(2)}(z_o)}{(q^{(7)}(z_o))^5} + 480 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^3 \frac{1}{120} q^{(10)}(z_o) p(z_o)}{(q^{(7)}(z_o))^5} \\
 & + 1440 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^3 \frac{1}{36} q^{(9)}(z_o) p^{(1)}(z_o)}{(q^{(7)}(z_o))^5} + \\
 & 1080 \times \frac{\left(\frac{1}{36} q^{(9)}(z_o)\right)^2 \left(\frac{1}{8} q^{(8)}(z_o)\right)^2 p(z_o)}{(q^{(7)}(z_o))^5} - 1800 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^4 \frac{1}{36} q^{(9)}(z_o) p(z_o)}{(q^{(7)}(z_o))^6} - \\
 & \left. 720 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^5 p^{(1)}(z_o)}{(q^{(7)}(z_o))^6} + 720 \times \frac{\left(\frac{1}{8} q^{(8)}(z_o)\right)^6 p(z_o)}{(q^{(7)}(z_o))^7} \right] \dots(22)
 \end{aligned}$$

Proof:

If $(z = z_o)$ is a pole of order $(\mathbf{m=7})$ then by (short – cut -method) we get :

$$\text{Res}(f, z_o) = \frac{1}{6!} \lim_{z \rightarrow z_o} \frac{d^6}{dz^6} \left[(z - z_o)^7 \frac{p(z)}{q(z)} \right] \text{ where } (f(z) = \frac{p(z)}{q(z)}) \dots(23)$$

We expand the analytic function $q(z)$ into Taylor series valid in disk $|z - z_o| < r$

$$= \frac{1}{6!} \lim_{z \rightarrow z_o} \frac{d^6}{dz^6} \left[\frac{(z - z_o)^7 p(z)}{q(z_o) + (z - z_o) q^{(1)}(z_o) + \frac{(z - z_o)^2}{2!} q^{(2)}(z_o) + \frac{(z - z_o)^3}{3!} q^{(3)}(z_o) + \dots} \right] \dots(24)$$

$$= 7 \lim_{z \rightarrow z_0} \frac{d^6}{dz^6} \left[\frac{p(z)}{U(z)} \right] \quad \dots(25)$$

where :

$$U(z) = q^{(7)}(z_0) + \frac{(z-z_0)}{8} q^{(8)}(z_0) + \frac{(z-z_0)^2}{8 \times 9} q^{(9)}(z_0) + \frac{(z-z_0)^3}{8 \times 9 \times 10} q^{(10)}(z_0) + \dots(26)$$

Then from (25) we get :

$$= 7 \lim_{z \rightarrow z_0} \frac{d^5}{dz^5} \left[\frac{U(z)p'(z) - p(z)U'(z)}{(U(z))^2} \right] \quad \dots(27)$$

$$= 7 \lim_{z \rightarrow z_0} \frac{d^4}{dz^4} \left[\frac{p^{(2)}(z)}{U(z)} - \frac{p(z)U^{(2)}(z)}{(U(z))^2} - 2 \times \frac{p^{(1)}(z)U'(z)}{(U(z))^2} + 2 \times \frac{p(z)U'(z)^2}{(U(z))^3} \right] \quad \dots(28)$$

$$= 7 \lim_{z \rightarrow z_0} \frac{d^3}{dz^3} \left[\frac{p^{(3)}(z)}{U(z)} - 3 \times \frac{p^{(2)}(z)U'(z)}{(U(z))^2} - \frac{p(z)U^{(3)}(z)}{(U(z))^2} - 3 \times \frac{p(z)U^{(2)}(z)}{(U(z))^2} \right. \\ \left. + 6 \times \frac{p(z)U'(z)U^{(2)}(z)}{(U(z))^3} + 6 \times \frac{p^{(1)}(z)U'(z)^2}{(U(z))^3} - 6 \times \frac{p(z)U'(z)^3}{(U(z))^4} \right] \quad \dots(29)$$

$$= 7 \lim_{z \rightarrow z_0} \frac{d^2}{dz^2} \left[\frac{p^{(4)}(z)}{U(z)} - 4 \times \frac{p^{(3)}(z)U'(z)}{(U(z))^2} \right. \\ - 6 \times \frac{p^{(2)}(z)U^{(2)}(z)}{(U(z))^2} + 12 \times \frac{p^{(2)}(z)U'(z)^2}{(U(z))^3} \\ - \frac{p(z)U^{(4)}(z)}{(U(z))^2} - 4 \times \frac{p^{(1)}(z)U^{(3)}(z)}{(U(z))^2} + 8 \times \frac{p(z)U'(z)U^{(3)}(z)}{(U(z))^3} \\ \left. + 24 \times \frac{p^{(1)}(z)U'(z)U^{(2)}(z)}{(U(z))^3} + \right. \\ \left. 6 \times \frac{p(z)U^{(2)}(z)^2}{(U(z))^3} - 36 \times \frac{p(z)U'(z)^2 U^{(2)}(z)}{(U(z))^4} \right. \\ \left. - 24 \times \frac{p^{(1)}(z)U'(z)^3}{(U(z))^4} + 24 \frac{p(z)U'(z)^4}{(U(z))^5} \right] \quad \dots(30)$$

$$= 7 \lim_{z \rightarrow z_0} \frac{d}{dz} \left[\frac{p^{(5)}(z)}{U(z)} - 5 \times \frac{p^{(4)}(z)U'(z)}{(U(z))^2} - 4 \times \frac{p^{(3)}(z)U^{(2)}(z)}{(U(z))^2} - 10 \times \frac{p^{(2)}(z)U^{(3)}(z)}{(U(z))^2} - \right.$$

$$\begin{aligned}
 & 6 \times \frac{p^{(2)}(z)U^{(2)}(z)}{(U(z))^2} - 5 \times \frac{p^{(1)}(z)U^{(4)}(z)}{(U(z))^2} - \frac{p(z)U^{(5)}(z)}{(U(z))^2} + 20 \times \frac{(U^{(1)}(z))^2 p^{(3)}(z)}{(U(z))^3} \\
 & + 60 \times \frac{U^{(1)}(z)U^{(2)}(z)p^{(2)}(z)}{(U(z))^3} + 10 \times \frac{p(z)U^{(1)}(z)U^{(4)}(z)}{(U(z))^3} + 8 \times \frac{U^{(1)}(z)U^{(2)}(z)p^{(1)}(z)}{(U(z))^3} \\
 & + 20 \times \frac{U^{(3)}(z)U^{(2)}(z)p(z)}{(U(z))^3} + 32 \times \frac{U^{(1)}(z)U^{(3)}(z)p^{(1)}(z)}{(U(z))^3} + 24 \times \frac{p^{(1)}(z)U^{(1)}(z)U^{(3)}(z)}{(U(z))^3} \\
 & + 30 \times \frac{p^{(1)}(z)(U^{(2)}(z))^2}{(U(z))^3} - 60 \times \frac{(U^{(1)}(z))^3 p^{(2)}(z)}{(U(z))^4} - 60 \times \frac{(U^{(1)}(z))^2 U^{(3)}(z)p(z)}{(U(z))^4} \\
 & - 180 \times \frac{(U^{(1)}(z))^2 U^{(2)}(z)p^{(1)}(z)}{(U(z))^4} - 90 \times \frac{U^{(1)}(z)(U^{(2)}(z))^2 p(z)}{(U(z))^4} + 240 \times \frac{(U^{(1)}(z))^3 U^{(2)}(z)p(z)}{(U(z))^5} \\
 & + 120 \times \frac{(U^{(1)}(z))^4 p^{(1)}(z)}{(U(z))^5} - 120 \times \frac{(U^{(1)}(z))^5 p(z)}{(U(z))^6} \quad \dots(31)
 \end{aligned}$$

$$\begin{aligned}
 & = 7 \lim_{z \rightarrow z_0} \left[\frac{p^{(6)}(z)}{U(z)} - 6 \times \frac{p^{(5)}(z)U^{(1)}(z)}{(U(z))^2} - 9 \times \frac{p^{(4)}(z)U^{(2)}(z)}{(U(z))^2} - 14 \times \frac{p^{(3)}(z)U^{(3)}(z)}{(U(z))^2} - \right. \\
 & 15 \times \frac{U^{(4)}(z)p^{(2)}(z)}{(U(z))^2} - 6 \times \frac{p^{(2)}(z)U^{(3)}(z)}{(U(z))^2} - 6 \times \frac{p^{(3)}(z)U^{(2)}(z)}{(U(z))^2} + 20 \times \frac{U^{(1)}(z)U^{(2)}(z)p^{(2)}(z)}{(U(z))^3} \\
 & - 6 \times \frac{p^{(1)}(z)U^{(5)}(z)}{(U(z))^2} - \frac{p(z)U^{(6)}(z)}{(U(z))^2} + 30 \times \frac{(U^{(1)}(z))^2 p^{(4)}(z)}{(U(z))^3} + 108 \times \frac{U^{(1)}(z)U^{(2)}(z)p^{(3)}(z)}{(U(z))^3} \\
 & + 80 \times \frac{U^{(1)}(z)U^{(3)}(z)p^{(2)}(z)}{(U(z))^3} + 76 \times \frac{U^{(1)}(z)U^{(4)}(z)p^{(1)}(z)}{(U(z))^3} + 12 \times \frac{U^{(1)}(z)U^{(5)}(z)p(z)}{(U(z))^3} \\
 & + 90 \times \frac{p^{(2)}(z)(U^{(2)}(z))^2}{(U(z))^3} + 30 \times \frac{p(z)U^{(2)}(z)U^{(4)}(z)}{(U(z))^3} + 8 \times \frac{U^{(1)}(z)U^{(3)}(z)p^{(1)}(z)}{(U(z))^3} \\
 & \left. + 8 \times \frac{p^{(1)}(z)(U^{(2)}(z))^2}{(U(z))^3} \right]
 \end{aligned}$$

$$\begin{aligned}
 & +136 \times \frac{U^{(2)}(z)U^{(3)}(z)p^{(1)}(z)}{(U(z))^3} + 20 \times \frac{(U^{(3)}(z))^2 p(z)}{(U(z))^3} \\
 & + 56 \times \frac{U^{(1)}(z)U^{(3)}(z)p^{(2)}(z)}{(U(z))^3} - 120 \times \frac{(U^{(1)}(z))^3 p^{(3)}(z)}{(U(z))^4} \\
 & - 540 \times \frac{(U^{(1)}(z))^2 U^{(2)}(z)p^{(2)}(z)}{(U(z))^4} - 90 \times \frac{(U^{(1)}(z))^2 U^{(4)}(z)p(z)}{(U(z))^4} \\
 & - 24 \times \frac{(U^{(1)}(z))^2 U^{(2)}(z)p^{(1)}(z)}{(U(z))^4} - \\
 & 360 \times \frac{U^{(1)}(z)U^{(2)}(z)U^{(3)}(z)p(z)}{(U(z))^4} + 408 \times \frac{(U^{(1)}(z))^2 U^{(3)}(z)p^{(1)}(z)}{(U(z))^4} \\
 & - 540 \times \frac{U^{(1)}(z)(U^{(2)}(z))^2 p^{(1)}(z)}{(U(z))^4} - 90 \times \frac{(U^{(2)}(z))^2 p(z)}{(U(z))^4} \\
 & + 360 \times \frac{(U^{(1)}(z))^4 p^{(2)}(z)}{(U(z))^5} + 480 \times \frac{(U^{(1)}(z))^4 U^{(3)}(z)p(z)}{(U(z))^5} + \\
 & 1440 \times \frac{(U^{(1)}(z))^3 U^{(2)}(z)p^{(1)}(z)}{(U(z))^5} + 1080 \times \frac{(U^{(2)}(z))^2 (U^{(1)}(z))^2 p(z)}{(U(z))^5} \\
 & - 1800 \times \frac{(U^{(1)}(z))^4 U^{(2)}(z)p(z)}{(U(z))^6} - 720 \times \frac{(U^{(1)}(z))^5 p^{(1)}(z)}{(U(z))^6} + 720 \times \frac{(U^{(1)}(z))^6 p(z)}{(U(z))^7}]
 \end{aligned} \tag{32}$$

Where :

$$\begin{aligned}
 U(z) &= q^{(7)}(z_0) + \frac{(z-z_0)}{8}q^{(8)}(z_0) + \frac{(z-z_0)^2}{9 \times 8}q^{(9)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U(z) &= q^{(7)}(z_0) \\
 U^{(1)}(z) &= \frac{1}{8}q^{(8)}(z_0) + \frac{2}{9 \times 8}(z-z_0)q^{(9)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U^{(1)}(z) &= \frac{1}{8}q^{(8)}(z_0) \\
 U^{(2)}(z) &= \frac{2}{9 \times 8}q^{(9)}(z_0) + \frac{3 \times 2}{10 \times 9 \times 8}(z-z_0)q^{(10)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U^{(2)}(z) &= \frac{1}{36}q^{(9)}(z_0) \\
 U^{(3)}(z) &= \frac{3 \times 2}{10 \times 9 \times 8}q^{(10)}(z_0) + \frac{4 \times 3 \times 2}{11 \times 10 \times 9 \times 8}(z-z_0)q^{(11)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U^{(3)}(z) &= \frac{1}{120}q^{(10)}(z_0) \\
 U^{(4)}(z) &= \frac{4 \times 3 \times 2}{11 \times 10 \times 9 \times 8}q^{(11)}(z_0) + \frac{5 \times 4 \times 3 \times 2}{12 \times 11 \times 10 \times 9 \times 8}(z-z_0)q^{(12)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U^{(4)}(z) &= \frac{1}{330}q^{(11)}(z_0) \\
 U^{(5)}(z) &= \frac{5 \times 4 \times 3 \times 2}{12 \times 11 \times 10 \times 9 \times 8}q^{(12)}(z_0) + \frac{6 \times 5 \times 4 \times 3 \times 2}{13 \times 12 \times 11 \times 10 \times 9 \times 8}(z-z_0)q^{(13)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U^{(5)}(z) &= \frac{1}{792}q^{(12)}(z_0) \\
 U^{(6)}(z) &= \frac{6 \times 5 \times 4 \times 3 \times 2}{13 \times 12 \times 11 \times 10 \times 9 \times 8}q^{(13)}(z_0) + \frac{7 \times 6 \times 5 \times 4 \times 3 \times 2}{14 \times 13 \times 12 \times 11 \times 10 \times 9 \times 8}(z-z_0)q^{(14)}(z_0) + \dots \\
 \lim_{z \rightarrow z_0} U^{(6)}(z) &= \frac{1}{1716}q^{(13)}(z_0)
 \end{aligned}
 \tag{33}$$

Then substituting the above values in (33) in equation (32) we get the desired proof .

3.6) In the above manner the procedure can be easily extended for any pole of order (m) .

4-Numerical computations :

These examples can be found in [1, 4, 6, 7, 8, 10, 11].

Example (4.1) :

Evaluate the following integral $\int_C \frac{Z^2 + 1}{(Z - 2)^4 (Z - 1)^3} dZ$ at $C : |Z| = 3$.

Solution:

The function $f(Z)$ has two poles ($Z=1, Z=2$) in the region $c:|Z|=3$, and the order of these poles is ($m=3$) & ($m=4$), respectively

The solution by Cauchy integral formula is as follows :

$$\int_{c:|Z|=3} \frac{Z^2 + 1}{(Z-2)^4(Z-1)^3} dZ = 58\pi i - 58\pi i = 0$$




Figure 4.1

Then the solution by the new procedure is given as follows :

$$\int_{c:|Z|=3} \frac{Z^2 + 1}{(Z-2)^4(Z-1)^3} dZ = 2\pi i \sum_{i=1}^2 \text{Res}(f, Z_i) = 2\pi i \text{Res}(f, Z_1 = 1) + 2\pi i \text{Res}(f, Z_2 = 2)$$

Where $p(Z) = Z^2 + 1$; $q(Z) = (Z-2)^4(Z-1)^3$

$P(2)=5$	$q^{(4)}(2) = 24$	$P(1)=2$	$q^{(3)}(1) = 6$
$p^{(1)}(2) = 4$	$q^{(5)}(2) = 360$	$p^{(1)}(1) = 2$	$q^{(4)}(1) = -96$
$p^{(2)}(2) = 2$	$q^{(6)}(2) = 2160$	$p^{(2)}(1) = 2$	$q^{(5)}(1) = 720$
$p^{(3)}(2) = 0$	$q^{(7)}(2) = 5040$	$p^{(3)}(3) = 0$	

Table 4.1

Then by using the new procedure for ($m=3$) we get :

$$\text{Res}(f,1) = 3 \times \left[\frac{1}{3} - 4 + 2.66 + \frac{2304}{(36 \times 6)} \right] = 28.98 \cong 29$$

Then by using the new procedure for ($m=4$) we get

$$\text{Res}(f,2) = 4 \times [-0.75 - 1.25 - 3 + 22.5 + 9 - 33.75] = -29$$

Then :

$$\int_{c:|Z|=3} \frac{Z^2 + 1}{(Z-2)^4(Z-1)^3} dZ = 2\pi i \sum_{i=1}^2 \text{Res}(f, Z_i) = 2\pi i \text{Res}(f, Z_1 = 1) + 2\pi i \text{Res}(f, Z_2 = 2) = 58\pi i - 58\pi i \cong 0$$

Example (4.2):

Evaluate the following integral $\int_c \frac{e^z - Z^6}{(Z-2i)^6} dz$ at $C:|Z|=3$.

Solution:

the function $f(z)$ has a pole $Z=2i$ of order ($m=6$).

then by Cauchy integral formula we get:

$$\int_{C:|Z|=3} \frac{e^Z - Z^6}{(Z - 2i)^6} dz = \frac{\pi i}{60} (e^{2i} - 1440i).$$

Then by using the new procedure for(m=6) we get

$$\int_{C:|Z|=3} \frac{e^z - Z^6}{(z - 2i)^6} dz = 2\pi i \operatorname{Res}(f, 2i) = \frac{\pi i}{60} (e^{2i} - 1440i).$$

Example (4.3)

Evaluate the following improper integral $\int_0^\infty \frac{\sin(x)}{x^m} dx$ where $m=1,3,5,\dots$

Solution:

Now for this type of improper integrations we not can find the result by Cauchy integral formula. The only way to evaluate is the new procedure with general order (m).

If (**m=1**) then $\int_0^\infty \frac{\sin(x)}{x} dx = \frac{\pi}{2}$

If (**m=3**) then $\int_0^\infty \frac{\sin(x)}{x^3} dx = \frac{-\pi}{4}$

If (**m=5**) then $\int_0^\infty \frac{\sin(x)}{x^5} dx = \frac{\pi}{48}$ we shall prove that




Figure 4.2

The function $f(Z) = \frac{e^Z}{Z^5}$ has pole ($Z=0$) of order ($m=5$), on real line. Then

$$\int_0^\infty \frac{\sin(x)}{x^5} dx = \lim_{R \rightarrow \infty} \int_0^R \frac{\sin(x)}{x^5} dx \quad \text{then by Jordan lemma we get:}$$

$$\lim_{R \rightarrow \infty, r \rightarrow 0} \left[\left(\int_{-R}^{-r} + \int_r^R + \int_{Sr} + \int_{CR} \right) \frac{\sin(x)}{x^5} dx \right] = 0$$

Where

$$\int \frac{\sin(x)}{x^5} dx = \int \frac{e^{iZ}}{Z^5} dZ \quad \text{on real part (x-axis). Then by lemma we get}$$

$$\left(\int_{CR} \frac{e^{iZ}}{Z^5} dz = 0 \right) \text{ then}$$

By replacing ($x = -x$) in the above first term in the LHS then we have :

$$\lim_{R \rightarrow \infty, r \rightarrow 0} \left[2i \int_r^R \frac{\sin(x)}{x^5} dx \right] = - \int_{Sr}^{iZ} \frac{e^{iZ}}{Z^5} dz = \pi i \operatorname{Res}(f, 0) \text{ then we find the residue at a}$$

pole ($z=0$) of order ($m=5$).

Using the new procedure to find the residue for order ($m=5$)

Now $p(z)=e^{iz}$ & $q(z)=Z^5$

$P(0)=1$	$q(0)=0$	$q^{(5)}(0) = 120$
$p^{(1)}(0) = i$	$q^{(1)}(0) = 0$	$q^{(6)}(0) = 0$
$p^{(2)}(0) = -1$	$q^{(2)}(0) = 0$	
$p^{(3)}(0) = -i$	$q^{(3)}(0) = 0$	
$p^{(4)}(0) = 1$	$q^{(4)}(0) = 0$	

Table 4.2

Then :

$$\operatorname{Res}(f, 0) = 5 \times \left[\frac{1}{120} \right] = \frac{1}{24} \text{ we get from that } \int_0^{\infty} \frac{\sin(x)}{x^5} dx = \frac{\pi}{48}$$

Then by using the new procedure of order (m) only we find general solution.

$$\int_0^{\infty} \frac{\sin(x)}{x^m} dx = \begin{cases} (-1)^{n+1} \frac{\pi}{2(m-1)!} & \text{if } m = 1,3,5,7,9,\dots \& n = 1,2,3,\dots \\ 0 & \text{otherwise} \end{cases}$$

Example (4.4)

Evaluate the following improper integral $\int_0^{\infty} \frac{\cos(x)}{x^m} dx$ Where $m=2,4,6,\dots$

Solution:

By the same way of example(3) we get the general solution .

$$\int_0^{\infty} \frac{\cos(x)}{x^m} dx = \begin{cases} (-1)^n \frac{\pi}{2(m-1)!} & \text{if } m = 2,4,\dots \& n = 1,2,3,\dots \\ 0 & \text{otherwise} \end{cases}$$

By using the new procedure of order (m) only we get the above solution.

5-conclusions :

It is clear that the new procedure for calculating poles of any order (m) is a very effective rule in determining the values of improper integrals while in the most of complex variable methods fail to compute these integrations.

REFERENCES

- [1] Trott, M. , (1997) "Visualization of Riemann Surfaces of Algebraic functions", **Mathematica in Education and Research**, Vol.(6), no. 4, pp.3-15.
- [2] Churchill, R. , (1974)" Complex Variable and Applications ", New York .
- [3] Rinehart and Winston , (1976) " Elements of Complex Variable " ,Macmilan & Co LTD London.
- [4] Dauglas, N ., (2000)" complex Analysis " .
- [5] Murray, R. , (1981) " Theory and problems of Complex Variable", Ginn & Company, Boston.
- [6] Oliver, K. , (1996) " The Residue theorem and it's Application_". New York .
- [7] Chen, L. (2003), " Introduction to complex analysis ".
- [8] Mike, P. (2005), " Complex Number and Complex Function ". New York .
- [9] Vitally, M. (2005)" Complex Function theory" , University of Bristol.
- [10] Sean, M. (2001) " Introduction to Applied mathematics ".
- [11] Risch, R., (1979) "Algebraic Properties of the Elementary Functions of Analysis", **American Journal of Mathematics**, Vol.(101), pp. 743-759.