

## Complex Dynamics of the family $\lambda \frac{\sinh^2(z)}{z^3}$

Salma M. Faris

College of Computer Sciences and Mathematics  
University of Mosul

Received on:12/5/2009

Accepted on:4/10/2009

### ABSTRACT

The aim of this work is to study the complex dynamics of the family  $F = \{ \lambda \frac{\sinh^2(z)}{z^3}, \lambda > 0 \}$ , of transcendental meromorphic functions. We prove that certain intervals are contained in  $J(f)$  or in  $F(f)$  and we prove that Julia set contains  $R^+ \cup iR^+$  and show that Fatou set of the functions in  $F$  contains certain components for different values of  $\lambda > 0$ . We characterize Julia set of  $f_\lambda \in F$ , for different values of  $\lambda > 0$ , as the closure of the escaping points, and use this characterization to give computer images for these sets.

**Keywords:** complex dynamics, Julia set, Fatou set.

الديناميكية المعقدة لعائلة الدوال  $\lambda \frac{\sinh^2(z)}{z^3}$

سلمى مصلح فارس

كلية علوم الحاسوب والرياضيات

جامعة الموصل

تاريخ القبول: 2009/10/4

تاريخ الاستلام: 2009/5/12

### الملخص

الهدف من البحث هو دراسة الدينامية المعقدة لعائلة الدوال الميرومورفية المتسامية  $F = \{ \lambda \frac{\sinh^2(z)}{z^3}, \lambda > 0 \}$ . سنبرهن أن فترات معينة من الأعداد الحقيقية تنتمي إلى مجموعة جوليا في حين تنتمي فترات أخرى إلى مجموعة فاتو كما سنبرهن على أن مجموعة جوليا تحوي على  $R^+ \cup iR^+$  كما سنبرهن أن مجموعة فاتو تحتوي على مكونات معينة ولمختلف قيم المعلمة  $\lambda > 0$ . بالإضافة إلى ذلك، سنعطي وصفاً لمجاميع جوليا على أنها أنغلاق مجموعة النقاط في الفضاء العقدي والتي تكون فيها غاية الدالة هو المالا نهائية لمختلف قيم المعلمة  $\lambda > 0$ . وسنستخدم هذا الوصف لإعطاء صور لهذه المجاميع بواسطة الحاسوب. الكلمات المفتاحية: الديناميكية المعقدة، مجموعة جوليا، مجموعة فاتو.

## 1. Introduction:

In the early of twentieth century, the iteration theory of complex functions originated in works of Julia and Fatou. There had been a long period of inactivity after that. During the end of 20-th century, a renewed interest in the study of iteration theory started due to beautiful computer graphics and a wide ranging applications in engineering and science associated with it [1,2,3,4,8, 10]. Iteration occurs in many parts of mathematics. For example many algorithms of numerical mathematics based on it. Given a set  $D$  and a function  $f : D \rightarrow D$  the iterates of  $f$  are defined by  $f^1 = f$  and  $f^n = f \circ f^{n-1}$  for  $n > 1$ . The main problem in iteration theory is to study the behavior of the sequence  $\langle f^n \rangle$  as  $n$  goes to infinity.

In iteration theory, for complex dynamical systems, most of the works have been centered around the dynamics of entire and rational functions. However, in comparison to the investigations of entire and rational functions, not much works have been done in this direction for transcendental meromorphic functions.

The initial works on study of transcendental meromorphic functions started at 80-th by Devaney, Keen, Bergweiler, and others [2,6,7]. The central objects studied in complex dynamical systems of a function is its Julia and Fatou sets. The Fatou set of a function  $f$ , denoted by  $F(f)$ , is defined to be the set of all complex numbers where the family of iterates  $\{f^n : n > 0\}$  of  $f$  forms a normal in the sense of Montel. Recall that the family  $F = \{f^n : n > 0\}$  of complex maps defined on open set  $U$  of  $C$  called normal if every infinite sequence in  $F$  contains a subsequence which either: converges uniformly on a compact subset of  $U$ , or converges uniformly to  $\infty$  on  $U$ [4].

## 2. Definitions:

Julia set (or chaotic set), denoted by  $J(f)$ , is the complement of  $F(f)$ . The escaping points set of meromorphic function  $f(z)$ , denoted by  $I(f)$ , is defined by  $I(f) = \{z \in C : f^{(n)}(z) \rightarrow \infty \text{ as } n \rightarrow \infty \text{ and } f^{(n)}(z) \neq \infty\}$ .

A point  $w \in C$  is said to be critical point of  $f$  if  $f'(z) = 0$ . The value  $f(w)$  corresponding to a critical point  $w$  is called a critical value of  $f$ . A point  $v \in C$  is called asymptotic value of  $f$  if there is a continuous function  $\alpha(t)$  satisfying that  $\lim_{t \rightarrow \infty} f(\alpha(t)) = v$ . A singular value of  $f$  is defined to be either critical value or asymptotic value of  $f$ . Finally the function  $f$  is called critically finite if it has a finite numbers of critical and asymptotic values.

### 3. Helping Results:

Let  $B$  be a class of meromorphic functions having bounded singular values. In this work, we need the following characterization given by Zheng [10], for transcendental meromorphic functions:

**Theorem (3.1):** Let  $f_\lambda \in B$ , then  $J(f) = \overline{I(f)}$ .

In [5], we studied the real dynamics of the family  $F = \lambda \frac{\sinh^2(z)}{z^3}$ . In present work we study the complex dynamics of this family. We show that the functions in this family are non-critically finite and all the critical values are bounded. We characterize the Julia set of  $f_\lambda \in F$ , as the escaping points of  $f_\lambda, \overline{I(f_\lambda)}$ , for various values of  $\lambda > 0$ . For different values of the parameter  $\lambda > 0$ , we prove that Julia and Fatou sets contains certain intervals. Finally we use the characterization of Julia set to give computer images of Julia set for different value of  $\lambda > 0$

In [5], it was shown that the critical parameter value of the family  $F$  is  $\lambda_1 = 1.2203$ , and we proved the following theorem :

**Theorem (3.2):** Let  $f_\lambda \in F$ . Let  $T_0$  be set of backward orbit of the pole  $z=0$ ,

then: **1.** For  $0 < \lambda < \lambda_1$ ,  $f_\lambda^n(x) \rightarrow a_\lambda$  for  $x \in (\alpha, r_\lambda)/T_0$ ,

$f_\lambda^n(x) \rightarrow -a_\lambda$  for  $x \in (-r_\lambda, -\alpha)/T_0$ . For  $x \in \{(0, \alpha) \cup (r_\lambda, \infty)\}$   $f_\lambda^n(x) \rightarrow \infty$

and, for  $x \in \{(-\alpha, 0) \cup (-\infty, -r_\lambda)\}$   $f_\lambda^n(x) \rightarrow -\infty$ , where  $\alpha$  is a positive solution of the equation  $f_\lambda(x) = r_\lambda$ ,  $a_\lambda$  is an attracting fixed point of  $f_\lambda(x)$  and  $r_\lambda$  is a repelling fixed point of  $f_\lambda(x)$ .

**2.** For  $\lambda = \lambda_1$ ,  $f_\lambda^n(x) \rightarrow x_1$  for  $x \in (\mu, x_1)/T_0$  and  $f_\lambda^n(x) \rightarrow -x$  for  $x \in (-x, -\mu)/T_0$

Moreover  $f_\lambda^n(x) \rightarrow \infty$  for

$x \in \{(0, \mu), (x_1, \infty)\}/T_0$  and  $f_\lambda^n(x) \rightarrow -\infty$  for  $x \in \{((-\infty, x_1) \cup (-\mu, 0))/T_0\}$ ,

where  $x_1$  and  $-x_1$  are the rationally indifferent fixed points of  $f_\lambda(x)$ . and  $\mu$  is a positive solution of the equation  $f_\lambda(x) = x_1$

**3. For**  $\lambda > \lambda_1$ ,  $f_\lambda^n(x) \rightarrow \infty$  for  $x \in (0, \infty)/T_0$  and  $f_\lambda^n(x) \rightarrow -\infty$  for  $x \in (-\infty, 0)/T_0$ .

The following theorem gives a classification of the periodic components of the Fatou set of a function  $f$ . Recall that a maximal connected domain  $U$  contained in the Fatou set of a function  $f$  is called a component of  $F(f)$ .

**Theorem (3.3)[9].**

Let  $f$  be a complex function. Let  $U$  be a periodic component, of period  $p$ , in the Fatou set of  $f$ . Then one of the following possibilities is true:

1.  $U$  contains an attracting periodic point  $z_0$  of period  $p$ . Then for  $z \in U$   $f^{(np)}(z) \rightarrow z_0$  as  $n \rightarrow \infty$  and  $U$  is called the **basin of attraction of  $z_0$** .
2.  $\partial U$  contains a periodic point  $z_0$ ,  $f^{(np)}(z) \rightarrow z_0$  as  $n \rightarrow \infty$  for  $z \in U$  and  $f^{(p)}(z_0) = 1$ . In this case  $U$  is called **Parabolic domain**.
3. There exists an analytic homeomorphism  $\varphi : U \rightarrow D$ , where  $D$  is the unit disk, such that  $\varphi \circ f^{(p)} \circ \varphi^{-1}(z) = e^{2\pi i \alpha} z$  for some  $\alpha \in R/Q$ . In this case  $U$  is called **Siegel disk**.
4. There exists an analytic homeomorphism  $\varphi : U \rightarrow A$ , where  $A = \{z \in C : 1 < |z| < r\}$  and  $r > 1$ , such that  $\varphi \circ f^{(p)} \circ \varphi^{-1}(z) = e^{2\pi i \alpha} z$  for some  $\alpha \in R/Q$ . In this case  $U$  is called **Herman ring**.
5. There exists  $z_0 \in U$  such that  $f^{(np)}(z) \rightarrow z_0$  as  $n \rightarrow \infty$  for  $z \in U$  but  $f^{(np)}(z_0)$  is not defined. In this case,  $U$  is called **Baker domain**.

If  $U$  is not periodic, then  $U$  is called **wandering domain**, i.e. if  $U$  is wandering domain, then  $U^m \neq U^n$  for all  $m \neq n$ .

**Theorem (3.4)[10]:** Let  $f$  be a transcendental meromorphic function and let  $D = \{U_0, U_1, \dots, U_{n-1}\}$  be a  $n$ -periodic cycle of components of  $F(f)$ . Then:

1. If  $D$  is a cycle of attracting basins or parabolic domains, then some  $U_k$ ,  $k=1, 2, \dots, n-1$ , must intersect the set of singular values.
2. if  $D$  is a Siegel disks or Herman rings, then  $U_k$  is a proper subset of the closure of the forward orbits of the singular values, for each  $k=1, 2, \dots, n-1$ .

**4. Main Results:**

In the following, we study the complex dynamics of the family  $F$  starting with the following proposition:

**Proposition (4.1):** Let  $f_\lambda \in F$ , then the function  $f_\lambda(z)$  is non-critically finite and all the singular values of it are bounded.

If we derive the function  $f_\lambda(z) = \lambda \frac{\sinh^2(z)}{z^3}$  and solve the equation  $f'_\lambda(z) = 0$ , then one can easily show that  $f_\lambda(z)$  has infinite number of critical values, two of them are real and the other are purely imaginary.

Moreover, if we assume that  $\{iy_n : n \in N\}$  is the sequence of imaginary critical value then  $\lambda \left| \frac{\sinh^2(iy)}{iy^3} \right| \leq \frac{\lambda}{|y_n^3|} < \lambda M$ , where  $M = \max\{1/y_n^3 : n \in N\}$ .

Therefore the function  $f_\lambda(z)$  is non-critically finite and all the singular values of it are bounded.

**Theorem (4.2):** Let  $f = f_\lambda \in F$ , then  $J(f) = \overline{I(f)}$ .

**Proof:** By proposition (4.1),  $f_\lambda \in B$ , where  $B$  is the class of meromorphic functions having bounded singular values. Then, by theorem (3.1),  $J(f) = \overline{I(f)}$ .

**Theorem (4.3):** let  $f_\lambda \in F$ . Then

1. For  $0 < \lambda < \lambda_1$ , the intervals  $\{(-r_\lambda, -\alpha) \cup (\alpha, r_\lambda)\}/T_0$  contained in  $F(f_\lambda)$ , and the intervals  $(0, \alpha) \cup (r_\lambda, \infty)$  are contained in  $J(f_\lambda)$ .
2. For  $\lambda = \lambda_1$ ,  $\{(-\mu, -x_1) \cup (x_1, \mu)\}/T_0$  contained in the Fatou set of  $f_\lambda$ ,  $F(f_\lambda)$ , and  $\{(0, \mu) \cup (x_0, \infty)\}$  are contained in the Julia set of  $f_\lambda$ ,  $J(f_\lambda)$ .
3. For  $\lambda > \lambda_1$ ,  $J(f_\lambda)$  contains  $R^+ \cup iR^+$ .

**Proof:**

1. For  $0 < \lambda < \lambda_1$ , by theorem (3.2), for  $x \in (\alpha, r_\lambda)/T_0$ ,  $f_\lambda^n(x) \rightarrow a_\lambda$  and  $x \in (-r_\lambda, -\alpha)/T_0$ ,  $f_\lambda^n(x) \rightarrow -a_\lambda$ . Thus for  $x \in (\alpha, r_\lambda)/T_0$ ,  $x$  is belong to the basin of attraction of the attracting fixed point  $a_\lambda$ ,  $B(a_\lambda)$ . But  $B(a_\lambda) \subseteq F(f_\lambda)$ . Thus  $(r_\lambda, \alpha)/T_0 \subseteq F(f_\lambda)$ . Similarly  $(-\alpha, -r_\lambda)/T_0 \subseteq B(-a_\lambda)$ , where  $B(-a_\lambda)$  is the basin of attraction of  $-a_\lambda$ . Hence  $(-\alpha, -r_\lambda)/T_0 \subseteq F(f_\lambda)$ . Therefore for  $0 < \lambda < \lambda_1$ ,  $\{(-r_\lambda, -\alpha) \cup (\alpha, r_\lambda)\}/T_0 \subseteq F(f_\lambda)$ . Also, by theorem (3.2) for  $x \in \{(0, \lambda) \cup (r_\lambda, \infty)\}/T_0$ ,  $f_\lambda^n(x) \rightarrow \infty$ . Hence, by theorem (3.3),  $\{(0, \lambda) \cup (r_\lambda, \infty)\}/T_0 \subseteq J(f_\lambda)$ . But the pole  $z=0$  and its pre images are contained in  $J(f_\lambda)$ . Thus  $\{(0, \lambda) \cup (r_\lambda, \infty)\}$  is contained in  $J(f_\lambda)$ .
2. For this case the proof is similar to that in part (1)
3. For  $\lambda > \lambda_1$ , by theorem (3.2),  $f_\lambda^n(x) \rightarrow \infty$  for  $x \in (0, \infty)/T_0$ . Thus  $(0, \infty)/T_0$  is contained in  $J(f_\lambda)$ . Since the pole  $z=0$  and its pre images are contained in  $J(f_\lambda)$ , then  $R^+$  is contained in  $J(f_\lambda)$ . But  $f_\lambda$  maps  $iR^+$  into

$R^+$ . Thus  $J(f_\lambda)$  contains the positive parts of the real and imaginary axes  $R^+ \cup iR^+$ .

**Theorem (4.4):** For  $f_\lambda \in F$ , we have the following:

1. For  $0 < \lambda < \lambda_1$ , the Fatou set,  $F(f_\lambda)$ , contains only two basins of attraction  $B(a_\lambda)$  and  $B(-a_\lambda)$ . Also,  $F(f_\lambda)$  has no any parabolic domains.
2. For  $\lambda = \lambda_1$ ,  $F(f_\lambda)$  has no parabolic domains other than that associated with the indifferent fixed points  $x_1$  and  $-x_1$ .
3. For  $\lambda > \lambda_1$ ,  $F(f_\lambda)$  has no basins of attraction, parabolic domains, Siegel disks or Herman rings.

**Proof:**

1. For  $0 < \lambda < \lambda_1$ . Let  $z \in B(a_\lambda)$ , by theorem (3.2),  $f_\lambda^n(z) \rightarrow a_\lambda$ . Hence the sequence  $\{f_\lambda^n(z)\}$  forms a normal family. Thus  $z \in F(f_\lambda)$ . Therefore  $B(a_\lambda) \subseteq F(f_\lambda)$ . Also, for any  $y \in B(-a_\lambda)$ , then  $f_\lambda^n(y) \rightarrow -a_\lambda$ . Hence  $\{f_\lambda^n(y)\}$  forms a normal family. Thus  $y \in F(f_\lambda)$ . Therefore  $B(-a_\lambda) \subseteq F(f_\lambda)$ . Again by theorem (3.2) part (a), the forward orbits of all singular values are either tend to  $a_\lambda$  or  $-a_\lambda$  or infinity. Hence by theorem (3.3),  $F(f_\lambda)$  has no other basins. Also, by theorem (3.3) does not contains any parabolic domains.

2. For  $\lambda = \lambda_1$ , by theorem (3.2),  $f_\lambda^n(x) \rightarrow x_1$  for  $x \in (\mu, x_1)/T_0$  and  $f_\lambda^n(x) \rightarrow \infty$  for  $x \in (x_1, \infty)/T_0$ .


Define  $U_1 = \{z \in C : f_\lambda^n(z) \rightarrow x_1 \text{ as } n \rightarrow \infty\}$ . Then the indifferent fixed points  $x_1$  belongs to the boundary of  $U_1$ . Hence  $U_1$  is a parabolic domains in  $F(f_\lambda)$ .

Similarly,  $U_2 = \{z \in C : f_\lambda^n(z) \rightarrow -x_1 \text{ as } n \rightarrow \infty\}$  is a parabolic domain of the indifferent fixed point  $-x_1$ . Again by theorem (3.2), the forward orbits of all singular values of  $f_\lambda$  tends to  $x_1$ ,  $-x_1$ , or infinity. Thus, using theorem (3.3),  $F(f_\lambda)$  has no basins of attraction and has no parabolic domains other than  $U_1$  and  $U_2$ .


3. For  $\lambda > \lambda_1$ , by theorem (3.2),  $f_\lambda^n(x) \rightarrow \infty$  for  $x \in (0, \infty) / T_0$ . Therefore the forward orbits of the critical values of  $f_\lambda$  are tend to infinity. Further, since the asymptotic value 0 is also a pole. Thus the orbit of 0 is tend to infinity. Hence,  $F(f_\lambda)$  has no any basins, parabolic domains, Siegel disks or Herman rings.

Finally we give computer images for Julia sets of the functions in the family  $F$  for  $0 < \lambda < \lambda_1$ ,  $\lambda = \lambda_1$  and  $\lambda > \lambda_1$  where  $\lambda_1 = 1.2203$  is the critical parameter value of the family  $F$ , see [5].


We used simple algorithm with Matlab 6.5 Program. A window has been selected in the plane and divided into  $k \times k$  grids with width  $d$ , we choose a large numbers  $N$  and  $M$ . For each grid point, compute the orbit up to maximum iteration  $N$ . If, at  $i < N$ , the modulus of the orbit is greater than  $M$ , then the grid point colored black and the iteration stopped. If not then original grid point left white. The output generated by this algorithm black and white pictures. The black points represent the approximated Julia set.



Julia set of the function  $\lambda \frac{\sinh^2(z)}{z^3}$  for  $\lambda < \lambda_1$



Julia set of the function  $\lambda \frac{\sinh^2(z)}{z^3}$  for  $\lambda = \lambda_1$



Julia set of the function  $\lambda \frac{\sinh^2(z)}{z^3}$  for  $\lambda > \lambda_1$

**REFERENCES**

- [1] Avitision A.A., Arakelion N.U., and A.A.Gonchar. **Asymptotic properties of meromorphic functions.** Akademii Nauk Asmyamskay SSR Seriya Matematika. 1989. V.24. no.3. pp. 207-222.
- [2] Barnsley M. F. & Stephen G. Demko., **Chaotic Dynamics and Fractals.** Academic Press. 1986.
- [3] Bruce J.. West & Bill Deering. **The rule of modern Science: Fractal Thinking. Studies of nonlinear phenomena un life sciences.** V. 3. J. World scientific, Singapore. 1995.
- [4] Devaney R. L. ., **Film and video as a tool in mathematical research.** The mathematical Intelligencer. 1989.
- [5] Faris S.M., **Real Dynamics and bifurcations of the family**  
 $F = \{\lambda \frac{\sinh^2(z)}{z^3} : \lambda > 0\}$  :To appear.
- [6] Gerald Tessch., **Differential Equations & Dynamical systems,** Fakullat of Math., Nordbegergstrab 15, 1090 Wien, Asttria, 2009.
- [7] Kapoor G.P.Prasad M.G.P. , **Dynamics of  $(e^z-1)/z$ : Julia set and Bifurcations,** Ergodic theory & Dyn.sys., vol.181, ( p.1362-1383), 1989.
- [8] Masahiro Nakagawa., **Chaos and Fractals in engineering.** World Scientific Pub. Co. 1999.
- [9] Nayak T. & Prasad M.G.P, **Bifurcation and Chaotic burst in the dynamics of  $\lambda e^{bz+\exp(z)}$**  , IITK, Guwahati,781093, India 2000.
- [10] Zheng J.H., **On transcendental meromorphic functions which are geometrically finite** , J. Ausrtral. Math. Soc. Vol.72, p. 93-170, 2002.